

NATIONAL GENERAL PRACTICE PROFILES

PROFILE FOR

SOUTHOWRAM SURGERY

LAW LANE, SOUTHOWRAM, HALIFAX





These profiles are designed to support GPs, clinical commissioning groups (CCGs) and PCTs to ensure that they are providing and commissioning effective and appropriate healthcare services for their local population. Using a variety of graphical displays such as spine charts and population pyramids, the tool presents a range of practice-level indicators drawn from the latest available data, including:

- local demography;
- Quality and Outcomes Framework domains;
- disease prevalence estimates;
- admission rates; and
- patient satisfaction.

In addition to displaying individual practice profiles, the web tool allows you to view summary profiles for PCTs and CCGs. Each practice can be compared with its PCT, CCG and with England, and also with the practices in the same deprivation decile and 'peer group' (although PDF generation is limited to PCT, CCG and England comparisons). The profiles do not provide an exhaustive list of primary care indicators, but they do allow a consistent approach to comparing and benchmarking across England. More indicators will be incorporated as the tool is developed further.

The profiles have been designed as a web tool and the full functionality is only available via the web version. For more information consult the User guide and FAQs via the Supporting documents link and for full metadata view 'Definitions' on the web pages.

The development of this tool has been led by erpho, the East of England Public Health Observatory, on behalf of the network of PHOs in England. For further information contact: feedback@erpho.org.uk



www.apho.org.uk/PracProf

How to read the indicator spine charts

The light grey bar shows the range of values found in England. The dark grey sections mark out the range within which the middle half of the observed values lie (25th to 75th percentile).

The red line shows where the England average is. The position of the circle shows the practice value, a diamond the PCT value, a triangle the cluster value, in relation to this scale.

The corresponding numbers can be found in the cells next to the chart.

If significance has been calculated for the indicator, then it is determined by whether the practice value is significantly higher or lower than the England average usually using 99.8% confidence intervals.

A significant difference is marked in blue, non-significance in yellow.

- No significant difference from England average
- Significantly different from England average
- Significance not calculated

- Practice
- △ PCT
- ▽ Clinical Commissioning Group



Practice Summary

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
% aged 0 to 4 years	2012	7.4%	6.2%	6.0%	0.0%		19.2%
% aged 5 to 14 years	2012	12.3%	11.9%	11.1%	0.0%		27.8%
% aged under 18 years	2012	23.4%	21.8%	20.8%	0.0%		64.6%
% aged 65+ years	2012	14.1%	15.8%	16.3%	0.0%		97.7%
% aged 75+ years	2012	5.6%	7.1%	7.6%	0.0%		84.2%
% aged 85+ years	2012	1.8%	2.1%	2.2%	0.0%		51.7%
Deprivation score (IMD)	2012	17.3	23.2	21.5	2.9		68.4
IDACI (Income Deprivation Affecting Children)	2012	15.0%	21.2%	21.8%	1.0%		68.0%
IDAOPI (Income Deprivation Affecting Older People)	2012	18.0%	19.4%	18.1%	4.0%		80.0%
% who would recommend practice	2011/12	90.9%	83.2%	81.9%	28.6%		100%
% satisfied with phone access	2011/12	82.2%	77.7%	78.0%	15.2%		100%
% satisfied with opening hours	2011/12	79.6%	80.1%	81.0%	40.2%		100%
% who saw/spoke to nurse or GP same or next day	2011/12	56.5%	52.8%	50.5%	11.5%		100%
% reporting good overall experience of making appointment	2011/12	81.7%	77.7%	79.1%	29.4%		100%
% who know how to contact an out-of-hours GP service	2011/12	51.7%	51.2%	58.0%	10.9%		88.3%
% with a long-standing health condition	2011/12	48.2%	54.1%	53.1%	19.6%		100%
% with health-related problems in daily life	2011/12	45.1%	49.2%	48.7%	12.8%		100%
% with caring responsibility	2011/12	24.1%	17.8%	18.4%	0.0%		38.7%
Disability allowance claimants (per 1000)	Nov 2011	40.8	49.4	48.3	8.7		235.1
Nursing home patients	2010/11	-	0.5%	0.5%	0.0%		66.9%
Working status - Paid work or full-time education	2011/12	60.3%	59.1%	60.7%	0.0%		100%
Working status - Unemployed	2011/12	8.2%	6.3%	5.6%	0.0%		81.0%
Total QOF points	2011/12	99.7%	98.4%	96.9%	34.8%		100%

Modelled Disease Prevalence

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Estimated prevalence of CVD (all ages)	2011	8.4%	9.5%	9.5%	2.6%		19.4%
Estimated prevalence of CHD (all ages)	2011	3.9%	5.1%	4.7%	0.0%		12.9%
Ratio of recorded vs expected CHD prevalence	2010/11	0.951	0.796	0.723	0.00		2.72
Estimated prevalence of COPD (all ages)	2011	2.3%	2.8%	2.9%	0.62%		6.47%
Ratio of recorded vs expected COPD prevalence	2010/11	0.535	0.621	0.559	0.00		2.54
Estimated prevalence of hypertension (all ages)	2011	23.5%	25.5%	24.9%	4.2%		44.7%
Ratio of recorded vs expected hypertension prevalence	2010/11	0.543	0.514	0.543	0.01		1.61
Estimated prevalence of stroke (all ages)	2011	1.7%	2.1%	2.1%	0.16%		5.44%
Ratio of recorded vs expected stroke prevalence	2010/11	1.01	0.871	0.827	0.00		2.84

CVD - Coronary heart disease

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
CHD: QOF prevalence (all ages)	2011/12	3.5%	4.0%	3.4%	0.0%		39.1%
Estimated prevalence of CHD (all ages)	2011	3.9%	5.1%	4.7%	0.0%		12.9%
Ratio of recorded vs expected CHD prevalence	2010/11	0.951	0.796	0.723	0.00		2.72
Heart failure w LVD: QOF prevalence	2011/12	0.6%	0.4%	0.4%	0.00%		4.95%
Exception rate for CHD indicators	2011/12	4.7%	8.2%	9.4%	0.0%		46.7%
CHD emergency admissions (per 100 patients on register)	2010/11	4.40%	5.60%	7.10%	0.0%		100.0%
CHD elective admissions (per 100 patients on register)	2010/11	4.40%	3.10%	5.25%	0.0%		100.0%
CHD 13: Angina referred for specialist assessment	2011/12	100%	99.1%	98.2%	0.0%		100%
CHD 6: Last BP reading in last 15mths is <=150/90	2011/12	95.5%	89.8%	90.1%	16.7%		100%
CHD 8: Last total cholesterol is <=5mmol/l	2011/12	89.0%	81.2%	80.4%	20.0%		100%
CHD 9: Record that aspirin, APT or ACT is taken	2011/12	94.3%	92.5%	93.3%	0.0%		100%
CHD 10: Currently treated with beta blocker	2011/12	65.9%	72.8%	74.2%	0.0%		100%
CHD 14: History of MI: treated with ACE-I	2011/12	100%	89.2%	91.1%	0.0%		100%
CHD 12: CHD patients given flu immunisation 1 Sep - 31 Mar	2011/12	91.1%	92.2%	92.5%	0.0%		100%

CVD - Stroke and TIA

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Stroke: QOF prevalence (all ages)	2011/12	1.4%	1.9%	1.7%	0.0%		23.6%
Estimated prevalence of stroke (all ages)	2011	1.7%	2.1%	2.1%	0.16%		5.44%
Ratio of recorded vs expected stroke prevalence	2010/11	1.01	0.871	0.827	0.00		2.84
Exception rate for stroke indicators	2011/12	6.9%	6.5%	7.8%	0.0%		77.8%
Stroke 13: New patients referred for further investigation	2011/12	80.0%	86.5%	89.6%	0.0%		100%
Stroke 6: Last BP reading is 150/90 or less	2011/12	82.9%	89.2%	88.6%	0.0%		100%
Stroke 7: Total cholesterol recorded in last 15mths	2011/12	91.7%	91.7%	91.4%	0.0%		100%
Stroke 8: Last measured total cholesterol <=5mmol/l	2011/12	81.3%	80.1%	77.2%	0.0%		100%
Stroke 12: Record of aspirin, APT or ACT taken	2011/12	96.0%	92.7%	93.6%	0.0%		100%
Stroke 10: Influenza immunisation given 1 Sep-31 Mar	2011/12	100%	90.3%	90.0%	0.0%		100%

CVD - Heart failure and atrial fibrillation

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Heart Failure: QOF prevalence (all ages)	2011/12	0.8%	0.7%	0.7%	0.00%		9.55%
Heart failure w LVD: QOF prevalence	2011/12	0.6%	0.4%	0.4%	0.00%		4.95%
Exception rate for heart failure indicators	2011/12	10.5%	11.6%	13.0%	0.0%		100%
Atrial fibrillation: QOF prevalence	2011/12	1.2%	1.4%	1.5%	0.0%		16.0%
Exception rate for atrial fibrillation indicators	2011/12	4.3%	3.2%	3.6%	0.0%		57.1%
HF 2: Diagnosis conf. by ECG/specialist assessm.	2011/12	100%	94.4%	95.7%	0.0%		100%
HF 3: HF w LVD: treated with ACE-I or ARB	2011/12	100%	88.6%	89.3%	0.0%		100%
HF 4: Heart failure w LVD: treatment w ACE inh. or ARB, and beta-blocker	2011/12	61.5%	82.1%	83.9%	0.0%		100%
AF 4: diagnosed with ECG or by specialist	2011/12	100%	96.4%	95.8%	0.0%		100%
AF 3: treated w anti- coag./platelet therapy	2011/12	92.9%	93.4%	93.7%	0.0%		100%
























CVD - Risk factors for CVD

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Hypertension: QOF prevalence (all ages)	2011/12	12.8%	13.2%	13.6%	0.0%		60.1%
Estimated prevalence of hypertension (all ages)	2011	23.5%	25.5%	24.9%	4.2%		44.7%
Ratio of recorded vs expected hypertension prevalence	2010/11	0.543	0.514	0.543	0.01		1.61
Exception rate for hypertension indicators	2011/12	2.6%	2.0%	2.5%	0.0%		31.5%
Exception rate for smoking indicators	2011/12	0.5%	0.6%	0.7%	0.0%		16.5%
Exception rate for CVD primary prevention indicators	2011/12	4.2%	7.3%	8.2%	0.0%		71.4%
Obesity: QOF prevalence (16+)	2011/12	14.3%	11.0%	10.7%	0.0%		41.8%
BP 4: Record of BP in last 9mths	2011/12	95.4%	91.6%	91.0%	0.0%		100%
BP 5: Last (9mths) blood pressure <=150/90	2011/12	80.4%	79.7%	79.7%	0.0%		100%
PP 1: CV risk assessment for new hypertension cases	2011/12	89.5%	81.6%	80.0%	0.0%		100%
PP 2: Lifestyle advice for new hypertension cases	2011/12	92.0%	83.5%	81.5%	0.0%		100%
Dep 1: Depression case finding in CHD and/or diabetes patients	2011/12	91.3%	88.7%	88.6%	0.0%		100%
Smoking 3: status recorded in last 15mths (certain conditions)	2011/12	97.0%	96.1%	95.6%	13.6%		100%
Smoking 4: cessation advice/referral offered (certain conditions)	2011/12	95.3%	90.8%	92.9%	8.8%		100%

Diabetes

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Diabetes: QOF prevalence (17+)	2011/12	5.6%	5.6%	5.8%	0.0%		18.0%
Exception rate for diabetes indicators	2011/12	6.8%	5.8%	6.9%	0.0%		36.9%
Hypertension: QOF prevalence (all ages)	2011/12	12.8%	13.2%	13.6%	0.0%		60.1%
Exception rate for hypertension indicators	2011/12	2.6%	2.0%	2.5%	0.0%		31.5%
Exception rate for smoking indicators	2011/12	0.5%	0.6%	0.7%	0.0%		16.5%
Obesity: QOF prevalence (16+)	2011/12	14.3%	11.0%	10.7%	0.0%		41.8%
Diabetes admissions (per 1000)	2010/11	0.000	0.700	1.15	0.0		11.5
Ratio of recorded vs expected diabetes prevalence	2008/09	0.962	0.886	0.883	0.0		12.0
DM 2: Record of BMI in the last 15mths	2011/12	96.3%	94.2%	94.9%	38.1%		100%
DM 26: Last HbA1c is <=7.5% in last 15mths	2011/12	74.5%	69.4%	69.9%	0.0%		100%
DM 27: Last HbA1c is <=8% in last 15mths	2011/12	78.6%	78.1%	78.7%	0.0%		100%
DM 28: Last HbA1c is <=9% in last 15mths	2011/12	90.4%	87.9%	88.6%	39.1%		100%
DM 21: Retinal screening in last 15mths	2011/12	96.2%	92.3%	91.9%	0.0%		100%
DM 29: Record of peripheral pulses last 15mths	2011/12	93.3%	93.3%	89.6%	0.0%		100%
DM 10: Record of neuropathy test last 15mths	2011/12	94.3%	94.9%	91.9%	0.0%		100%
DM 30: Last BP is <=150/90	2011/12	91.7%	89.2%	89.9%	15.0%		100%
DM 31: Last BP is <=140/80	2011/12	74.3%	70.4%	70.7%	0.0%		100%
DM 13: Record of micro-albuminuria test last 15mths	2011/12	94.1%	90.0%	88.9%	0.0%		100%
DM 22: eGFR or serum creatinin testing in last 15mths	2011/12	99.1%	96.9%	96.9%	40.0%		100%
DM 15: Proteinuria/micro-album. treated w inhibitors	2011/12	92.3%	86.8%	87.4%	0.0%		100%
DM 17: Measured total chol (last 15mths) <=5mmol/l	2011/12	82.7%	83.9%	81.7%	0.0%		100%
DM 18: Influenza immunisation given 1 Sep - 31 Mar	2011/12	93.7%	91.4%	90.7%	32.0%		100%
Dep 1: Depression case finding in CHD and/or diabetes patients	2011/12	91.3%	88.7%	88.6%	0.0%		100%
Smoking 3: status recorded in last 15mths (certain conditions)	2011/12	97.0%	96.1%	95.6%	13.6%		100%
Smoking 4: cessation advice/referral offered (certain conditions)	2011/12	95.3%	90.8%	92.9%	8.8%		100%

Mental Health

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Psychoses: QOF prevalence (all ages)	2010/11	0.6%	0.8%	0.8%	0.0%		23.8%
Exception rate for patients with psychoses indicators	2011/12	6.4%	11.6%	11.8%	0.0%		53.9%
Dementia: QOF prevalence (all ages)	2011/12	0.4%	0.5%	0.5%	0.0%		58.7%
Exception rate for dementia indicators	2011/12	16.7%	11.5%	13.4%	0.0%		100%
Depression: QOF prevalence (18+)	2011/12	17.0%	14.9%	11.7%	0.0%		49.8%
Exception rate for depression indicators	2011/12	5.3%	5.6%	5.7%	0.0%		84.6%
% reporting a long-term mental health problem	2011/12	6.9%	4.6%	4.2%	0.0%		56.8%
MH 10: comprehensive care plan	2011/12	100%	91.1%	88.4%	0.0%		100%
MH 11: record of alcohol consumption	2011/12	92.9%	90.7%	89.6%	0.0%		100%
MH 12: record of BMI	2011/12	100%	90.2%	88.7%	0.0%		100%
MH 13: record of blood pressure	2011/12	100%	93.9%	91.7%	0.0%		100%
MH 14: record of cholesterol:hdl ratio	2011/12	80.0%	80.1%	81.9%	0.0%		100%
MH 15: record of blood glucose level	2011/12	88.9%	85.1%	84.8%	0.0%		100%
MH 16: record of cervical screening test	2011/12	100%	87.8%	88.0%	0.0%		100%
MH 7: Pychoses patients who DNA review: followed up < 14 d	2010/11	100%	96.7%	94.6%	0.0%		100%
MH 17: Li-therapy: record of s.creat. & TSH last 9mths	2011/12	100%	98.5%	96.3%	0.0%		100%
MH 18: Lithium therapy: record of Li-levels last 4mths	2011/12	100%	89.9%	89.3%	0.0%		100%
Dem 2: Dementia care has been reviewed last 15mths	2011/12	88.9%	78.0%	79.3%	0.0%		100%
DEM 3: Blood tests recorded	2011/12	100%	89.3%	83.5%	0.0%		100%
% reporting Alzheimer's disease or dementia	2011/12	1.8%	0.8%	0.6%	0.0%		60.0%
Dep 1: Depression case finding in CHD and/or diabetes patients	2011/12	91.3%	88.7%	88.6%	0.0%		100%
Dep 4: Depression cases with severity assessment	2011/12	93.9%	90.3%	91.4%	0.0%		100%
Dep 5: Depression cases with second severity assessment	2011/12	89.3%	72.5%	71.8%	0.0%		100%

Respiratory Disease

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
COPD: QOF prevalence (all ages)	2011/12	1.5%	1.8%	1.7%	0.0%		14.1%
Estimated prevalence of COPD (all ages)	2011	2.3%	2.8%	2.9%	0.62%		6.47%
Ratio of recorded vs expected COPD prevalence	2010/11	0.535	0.621	0.559	0.00		2.54
Exception rate for COPD indicators	2011/12	5.2%	10.2%	11.8%	0.0%		100%
COPD admissions (per 1000)	2010/11	3.30	2.60	2.12	0.0		18.5
Asthma: QOF prevalence (all ages)	2011/12	5.8%	6.3%	5.9%	0.1%		86.9%
Exception rate for asthma indicators	2011/12	2.1%	3.6%	5.3%	0.0%		52.1%
Exception rate for smoking indicators	2011/12	0.5%	0.6%	0.7%	0.0%		16.5%
COPD 15: Diagnosis conf. by spirometry	2011/12	100%	94.8%	93.0%	0.0%		100%
COPD 10: Record of FEV1 in last 15mths	2011/12	91.4%	90.0%	88.8%	0.0%		100%
COPD 13: assessed using MRC dyspnoea score last 15mths	2011/12	94.6%	91.3%	91.8%	0.0%		100%
COPD 8: Influenza immunisation given 1 Sep - 31 Mar	2011/12	94.1%	94.3%	93.1%	0.0%		100%
Asthma 8: with measures of variability/reversibility (8+)	2011/12	83.3%	87.9%	87.2%	0.0%		100%
Asthma 3: smoking recorded in last 15 mths (14-19y w asthma)	2011/12	83.3%	88.2%	89.0%	0.0%		100%
Asthma 6: review in the last 15mths	2011/12	90.3%	76.8%	78.1%	13.2%		100%
Smoking 3: status recorded in last 15mths (certain conditions)	2011/12	97.0%	96.1%	95.6%	13.6%		100%
Smoking 4: cessation advice/referral offered (certain conditions)	2011/12	95.3%	90.8%	92.9%	8.8%		100%
Uptake seasonal flu vaccine (65+)	2010/11	73.9%	70.7%	72.8%	9.0%		100%
Uptake seasonal influenza vaccine (6 months to 64 years old)	2010/11	54.2%	50.0%	50.4%	3.0%		100%

Chronic Kidney Disease

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
CKD: QOF prevalence (18+)	2011/12	4.5%	4.5%	4.3%	0.0%		40.5%
Exception rate for CKD indicators	2011/12	6.7%	3.0%	3.7%	0.0%		100%
CKD 2: Record of BP in last 15mths	2011/12	97.7%	98.0%	97.2%	17.1%		100%
CKD 3: Last BP reading measured in last 15mths is <=140/85	2011/12	78.6%	74.9%	75.1%	0.0%		100%
CKD 5: Hypertension treated with ACE inhibitor/ARB	2011/12	88.9%	89.2%	89.5%	0.0%		100%
CKD 6: Urine albumin:creatinine ratio test last 15mths	2011/12	90.9%	83.6%	82.2%	0.0%		100%

Other Conditions

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Cancer: QOF prevalence (all ages)	2011/12	1.5%	1.8%	1.8%	0.0%		19.1%
Exception rate for cancer indicators	2011/12	0.0%	0.8%	1.5%	0.0%		100%
Cancer admissions (per 1000)	2010/11	24.5	31.3	28.6	0.0		112.2
Cancer 3: review within 6mths of diagnosis	2011/12	100%	90.7%	93.3%	0.0%		100%
% reporting cancer in the last 5 years	2011/12	1.8%	3.0%	3.0%	0.0%		11.1%
Epilepsy: QOF prevalence (18+)	2011/12	0.8%	1.0%	0.8%	0.0%		12.9%
Exception rate for epilepsy indicators	2011/12	8.1%	14.3%	13.5%	0.0%		100%
Epilepsy 6: Record of seizure frequency	2011/12	100%	95.7%	95.2%	0.0%		100%
Epilepsy 8: Seizure free for last 12mths	2011/12	80.0%	72.5%	74.7%	0.0%		100%
Epilepsy 9: Contraception, conception and pregnancy	2011/12	100%	91.1%	90.0%	0.0%		100%
Learning disability: QOF prevalence (18+)	2011/12	0.2%	0.5%	0.5%	0.00%		5.90%
% reporting learning difficulty	2011/12	0.0%	0.8%	1.3%	0.0%		13.6%
LD 2: Down's Syndrome with blood TSH record	2011/12	-	90.5%	90.4%	0.0%		100%
Hypothyroidism: QOF prevalence (all ages)	2011/12	3.1%	4.1%	3.1%	0.0%		28.2%
Exception rate for hypothyroidism indicators	2011/12	0.0%	0.2%	0.5%	0.0%		44.4%
Thyroid 2: function test recorded last 15mths	2011/12	98.7%	96.7%	96.0%	11.4%		100%
Palliative/supportive care: QOF prevalence (all ages)	2011/12	0.3%	0.2%	0.2%	0.0%		31.7%
% reporting blindness or severe visual impairment	2011/12	0.7%	1.2%	1.1%	0.0%		11.0%
% reporting deafness or severe hearing impairment	2011/12	4.2%	4.3%	3.9%	0.0%		31.4%
% reporting a long-term back problem	2011/12	10.0%	9.1%	10.1%	0.0%		35.0%
% reporting arthritis or long-term joint problem	2011/12	17.0%	14.0%	13.1%	0.0%		53.7%

Secondary Care Use - Outpatients

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
All outpatient attendances (per 1000)	2010/11	801.1	792.8	852.5	138		2541
Total outpatient appointment expenditure (£ per patient)	2010	85.1	89.2	91.8	0.0		222.0
First outpatient appointment (per 1000)	2010	308.8	318.6	346.8	0		1077
First outpatient appointment expenditure (£ per patient)	2010	42.8	44.0	43.2	0.0		138.7
GP referrals to outpatients - 1st attendance (per 1000)	2010/11	153.1	171.6	191.7	29.6		557.2
Follow-ups to first outpatient appointments (per 1000)	2010	470.8	492.4	549.7	0		1384
Outpatients discharged at 1st appointment (per 1000)	2010	95.4	95.4	110.0	5.7		583.1
DNA: Rate of total outpatient appointments (per 1000)	2010	252.5	249.9	190.7	0.0		974.3
DNA: proportion of total 1st outpatient appointments	2010/11	6.1%	6.3%	7.0%	0.0%		41.4%
GP referrals to General Medicine - 1st attendance (per 1000)	2010/11	5.30	5.80	10.8	0.0		117.8
General Medicine - return ratio	2010/11	2.90	2.80	2.50	0.0		17.2
GP referrals to General Surgery - 1st attendance (per 1000)	2010/11	20.4	27.2	23.9	0.0		70.5
General Surgery - return ratio	2010/11	0.800	0.800	1.40	0.00		4.90
GP referrals to Gynaecology - 1st attendance (per 1000)	2010/11	16.7	14.3	17.1	0.0		80.2
Gynaecology - return ratio	2010/11	1.50	1.30	1.20	0.00		4.20
GP referrals to Orthopaedics - 1st attendance (per 1000)	2010/11	27.0	28.6	21.3	0.0		191.6
Orthopaedics - return ratio	2010/11	1.70	1.70	1.80	0.50		3.80
GP Referrals to Urology - 1st attendance (per 1000)	2010/11	7.80	8.70	7.97	0.0		27.4
Urology - return ratio	2010/11	2.30	2.00	2.20	0.0		15.0
GP referrals to Paediatrics - 1st attendance (per 1000)	2010/11	4.50	5.40	6.15	0.0		78.9
GP referrals to Dermatology - 1st attendance (per 1000)	2010/11	9.80	10.00	13.9	0.0		56.1

Secondary Care Use - A&E and Inpatients

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
A&E attendances (per 1000)	2010/11	294.0	310.2	387.2	14		1941
A&E admissions (per 1000)	2010/11	11.4	14.6	9.85	0.0		307.2
A&E attendances with primary diagnosis recorded	2010/11	2.8%	5.0%	59.8%	1.0%		100%
A&E referrals (weighted)	2010	4.06	7.69	16.8	0.0		398.4
Total admissions (per 1000)	2010/11	252.8	270.8	234.8	11		1005
Elective admissions (per 1000)	2010/11	115.2	125.7	121.0	6.0		327.2
Emergency admissions (per 1000)	2010/11	91.1	107.6	89.0	3.9		815.2
Elective inpatient admissions (per 1000)	2010/11	26.9	32.4	25.6	0.0		78.3
Day case admissions (per 1000)	2010/11	88.2	94.0	93.2	4.9		277.5
Long term conditions emergency bed days (per 1000)	2010/11	373.6	501.6	469.6	17		7247
Average emergency overnight occupied beds (per 1000)	2010/11	1.22	1.37	1.22	0.0		16.2
CHD total admissions (per 1000)	2010/11	4.10	4.90	4.57	0.0		24.1
CHD elective admissions (per 100 patients on register)	2010/11	4.40%	3.10%	5.25%	0.0%		100.0%
CHD emergency admissions (per 100 patients on register)	2010/11	4.40%	5.60%	7.10%	0.0%		100.0%
Respiratory disease admissions (per 1000)	2010/11	18.4	20.5	16.8	0.0		146.9
Diabetes admissions (per 1000)	2010/11	0.000	0.700	1.15	0.0		11.5
COPD admissions (per 1000)	2010/11	3.30	2.60	2.12	0.0		18.5
Cancer admissions (per 1000)	2010/11	24.5	31.3	28.6	0.0		112.2
Admissions for long term neurological conditions (per 1000)	2010/11	4.08	6.50	5.71	0.5		25.0
Emergency admissions for chronic conditions (per 1000)	2010/11	18.0	18.3	15.0	0.0		186.4
Total admissions expenditure (£ per patient)	2010	311.7	370.7	332.9	0		2161
Expenditure on elective admissions (£ per patient)	2010	127.1	141.7	137.2	0.0		318.5
Expenditure on emergency admissions (£ per patient)	2010	137.8	181.3	156.9	0		1880

Indicators for Needs Assessment

Indicator	Period	Practice Value	PCT Value	England Average	England Lowest	England Range	England Highest
Life expectancy - MSOA based estimate	2006 - 10	83.9	81.6	82.3	75.1		92.0
Fertility rate (MSOA based)	2006 - 10	66.1	66.9	63.1	12.0		134.3
Low birthweight births - MSOA based estimate	2006 - 10	8.3%	8.3%	7.5%	2.4%		13.0%
All cause elective standardised admission ratio (MSOA based)	2006/07 - 10/11	99.3	95.7	100.0	47.7		174.8
All cause emergency standardised admission ratio (MSOA based)	2006/07 - 10/11	107.9	107.6	100.0	43.8		235.2
CHD elective standardised admission ratio (MSOA based)	2006/07 - 10/11	70.5	60.1	100.0	32.6		293.5
CHD emergency standardised admission ratio (MSOA based)	2006/07 - 10/11	107.6	106.1	100.0	40.2		376.0
Stroke emergency standardised admission ratio (MSOA based)	2006/07 - 10/11	91.9	90.9	100.0	43.3		212.3
COPD emergency standardised admission ratio (MSOA based)	2006/07 - 10/11	140.4	131.0	100.0	17.7		541.8
Limiting long-term illness (MSOA based)	2009	17.8%	17.7%	16.9%	6.0%		32.0%
Long term unemployment (MSOA based)	2010/11	4.7	5.3	6.0	0.1		40.8
Healthy eating adults (MSOA based)	2006 - 08	24.1%	25.6%	28.7%	13.8%		54.1%
Prevalence of adults who binge drink	2007 - 08	23.1%	23.3%	20.0%	4.3%		50.4%
Standardised admission ratio for alcohol attributable harm (MSOA based)	2006/07 - 10/11	110.2	106.7	100.0	34.0		273.7
Prevalence of obese adults	2006 - 08	29.0%	27.0%	24.2%	9.6%		33.8%
Incidence of all cancers (MSOA based)	2005 - 09	112.5	102.0	100.0	66.7		147.0
Cancer mortality age <75 (MSOA based)	2006 - 10	103.3	106.2	100.0	37.2		205.3
Circulatory disease mortality age <75 (MSOA based)	2006 - 10	107.2	115.6	100.0	26.8		271.9